

Signatures of Universal Characteristics of Fractal Fluctuations in Global Mean Monthly Temperature Anomalies

A. M. Selvam

Deputy Director (Retired)

Indian Institute of Tropical Meteorology, Pune 411 008, India

Email: amselvam@gmail.com

Web sites: <http://www.geocities.com/amselvam>

<http://amselvam.tripod.com/index.html>

Running head: Universal characteristics of fractal fluctuations

Selfsimilar space-time fractal fluctuations are generic to dynamical systems in nature such as atmospheric flows, heartbeat patterns, population dynamics, etc. The physics of the long-range correlations intrinsic to fractal fluctuations is not yet identified. It is important to quantify the physics underlying the irregular fractal fluctuations for predictability studies. A general systems theory for fractals visualizes the emergence of successively larger scale fluctuations to result from the space-time integration of enclosed smaller scale fluctuations. The model predictions are as follows. The probability distribution and the power spectrum for fractal fluctuations is the same inverse power law function incorporating the *golden mean*. The predicted distribution is close to the Gaussian distribution for small-scale fluctuations but exhibits *fat long tail* for large-scale fluctuations with higher probability of occurrence than predicted by Gaussian distribution. Since the power (variance i.e. square of eddy amplitude) spectrum also represents the probability densities as in the case of quantum systems such as the electron or photon, fractal fluctuations exhibit quantumlike chaos. The *fine structure constant* for spectrum of fractal fluctuations is a function of the golden mean and is analogous to atomic spectra equal to about 1/137. Global gridded time series data sets of monthly mean temperatures for the period 1880 – 2007/2008 are analysed to show that data sets and corresponding power spectra exhibit distributions close to the model predicted inverse power law distribution. The model predicted and observed universal spectrum for interannual variability rules out linear secular trends in global monthly mean temperatures. Global warming results in intensification of fluctuations of all scales and manifested immediately in high frequency fluctuations.

Key words: fractals and statistical normal distribution, power law distributions, long-range correlations and fat tail distributions, golden mean and fractal fluctuations

Signatures of Universal Characteristics of Fractal Fluctuations in Global Monthly Mean Temperature Data

A. M. Selvam

Deputy Director (Retired)

Indian Institute of Tropical Meteorology, Pune 411 008, India

Email: amselvam@gmail.com

Web sites: <http://www.geocities.com/amselvam>

<http://amselvam.tripod.com/index.html>

Selfsimilar space-time fractal fluctuations are generic to dynamical systems in nature such as atmospheric flows, heartbeat patterns, population dynamics, etc. The physics of the long-range correlations intrinsic to fractal fluctuations is not yet identified. It is important to quantify the physics underlying the irregular fractal fluctuations for predictability studies. A general systems theory for fractals visualizes the emergence of successively larger scale fluctuations to result from the space-time integration of enclosed smaller scale fluctuations. The model predictions are as follows. The probability distribution and the power spectrum for fractal fluctuations is the same inverse power law function incorporating the *golden mean*. The predicted distribution is close to the Gaussian distribution for small-scale fluctuations but exhibits *fat long tail* for large-scale fluctuations with higher probability of occurrence than predicted by Gaussian distribution. Since the power (variance, i.e. square of eddy amplitude) spectrum also represents the probability densities as in the case of quantum systems such as the electron or photon, fractal fluctuations exhibit quantumlike chaos. The *fine structure constant* for spectrum of fractal fluctuations is a function of the golden mean and is analogous to atomic spectra equal to about 1/137. Global gridded time series data sets of monthly mean temperatures for the period 1880 – 2007/2008 are analysed to show that data sets and corresponding power spectra exhibit distributions close to the model predicted inverse power law distribution. The model predicted and observed universal spectrum for interannual variability rules out linear secular trends in global monthly mean temperatures. Global warming results in intensification of fluctuations of all scales and manifested immediately in high frequency fluctuations.

1. Introduction

Dynamical systems in nature such as atmospheric flows, heartbeat patterns, population dynamics, stock market indices, DNA base A, C, G, T sequence pattern, etc., exhibit irregular space-time fluctuations on all scales and exact quantification of the fluctuation pattern for predictability purposes has not yet been achieved. Traditionally, the Gaussian probability distribution is used for a broad quantification of the data set variability in terms of the sample mean and variance. The fractal or selfsimilar nature of space-time fluctuations was identified by Mandelbrot (1975) in the 1970s. Representative examples of fractal fluctuations of monthly mean temperatures for the period 1880 to 2007/2008 are shown Fig. 1.

fractal fluctuations
gridded temperature anomalies
Global Historical Climatology Network
(GHCN-Monthly) data Jan - Dec 1880-2007/2008




Fig. 1: Representative examples of fractal fluctuations of Global gridded monthly mean temperature anomalies for the period 1880 – 2007/2008

Fractal fluctuations (Fig. 1) show a zigzag selfsimilar pattern of successive increase followed by decrease on all scales (space-time), for example in atmospheric flows, cycles of increase and decrease in meteorological parameters such as wind, temperature, etc. occur from the turbulence scale of millimeters-seconds to climate scales of thousands of kilometers-years. The power spectra of fractal fluctuations exhibit inverse power law of the form f^α where f is the frequency and α is a constant. Inverse power law for power spectra indicate long-range space-time correlations or

scale invariance for the scale range for which α is a constant, i.e., the amplitudes of the eddy fluctuations in this scale range are a function of the scale factor α alone. In general the value of α is different for different scale ranges indicating multifractal structure for the fluctuations. The long-range space-time correlations exhibited by dynamical systems are identified as self-organized criticality (Bak *et al.*, 1988; Schroeder, 1990). The physics of self-organized criticality is not yet identified. The physics of fractal fluctuations generic to dynamical systems in nature is not yet identified and traditional statistical, mathematical theories do not provide adequate tools for identification and quantitative description of the observed universal properties of fractal structures observed in all fields of science and other areas of human interest. A recently developed general systems theory for fractal space-time fluctuations (Selvam, 1990, 2005, 2007; Selvam and Fadnavis, 1998) shows that the larger scale fluctuation can be visualized to emerge from the space-time averaging of enclosed small scale fluctuations, thereby generating a hierarchy of selfsimilar fluctuations manifested as the observed eddy continuum in power spectral analyses of fractal fluctuations. Such a concept results in inverse power law form incorporating the golden mean τ for the space-time fluctuation pattern and also for the power spectra of the fluctuations (Sec. 3). The predicted distribution is close to the Gaussian distribution for small-scale fluctuations, but exhibits *fat long tail* for large-scale fluctuations. Analyses of extensive data sets of Global gridded data sets of monthly mean temperatures for the period 1880 – 2007/2008 show that the space/time data sets follow closely, but not exactly the statistical normal distribution, particularly in the region of normalized deviations t greater than 2, the t values being computed as equal to $(x-av)/sd$ where av and sd denote respectively the mean and standard deviation of the variable x . The general systems theory, originally developed for turbulent fluid flows, provides universal quantification of physics underlying fractal fluctuations and is applicable to all dynamical systems in nature independent of its physical, chemical, electrical, or any other intrinsic characteristic. In the following, Sec. 2 gives a summary of traditional statistical and mathematical theories/techniques used for analysis and quantification of space-time fluctuation data sets. The general systems theory for fractal space-time fluctuations is described in Sec. 3. Sec. 4 deals with data and analyses techniques. Discussion and conclusions of results are presented in Sec. 5.

2. Statistical methods for data analysis

Dynamical systems such as atmospheric flows, stock markets, heartbeat patterns, population growth, traffic flows, etc., exhibit irregular space-time fluctuation patterns. Quantification of the space-time fluctuation pattern will help predictability studies, in particular for events which affect day-to-day human life such as extreme weather events, stock market crashes, traffic jams, etc. The analysis of data sets and broad quantification in terms of probabilities belongs to the field of statistics. Early attempts resulted in identification of the following two quantitative (mathematical) distributions which approximately fit data sets from a wide range of scientific and other disciplines of study. The first is the well known statistical normal distribution and the second is the power law distribution associated with the recently identified '*fractals*' or selfsimilar characteristic of data sets in general. In the following, a summary is given of the history and merits of the two distributions.

2.1 Statistical normal distribution

Historically, our present day methods of handling experimental data have their roots about four hundred years ago. At that time scientists began to calculate the odds in gambling games. From those studies emerged the theory of probability and subsequently the theory of statistics. These new statistical ideas suggested a different and more powerful experimental approach. The basic idea was that in some experiments random errors would make the value measured a bit higher and in other experiments random errors would make the value measured a bit lower. Combining these values by computing the average of the different experimental results would make the errors cancel and the average would be closer to the "right" value than the result of any one experiment (Liebovitch and Scheurle, 2000).

Abraham de Moivre, an 18th century statistician and consultant to gamblers made the first recorded discovery of the normal curve of error (or the bell curve because of its shape) in 1733. The normal distribution is the limiting case of the binomial distribution resulting from random operations such as flipping coins or rolling dice. Serious interest in the distribution of errors on the part of mathematicians such as Laplace and Gauss awaited the early nineteenth century when astronomers found the bell curve to be a useful tool to take into consideration the errors they made in their observations of the orbits of the planets (Goertzel and Fashing, 1981, 1986). The importance of the normal curve stems primarily from the fact that the distributions of many natural phenomena are at least approximately normally distributed. This normal distribution concept has molded how we analyze experimental data over the last two hundred years. We have come to think of data as having values most of which are near an average value, with a few values that are smaller, and a few that are larger. The probability density function $PDF(x)$ is the probability that any measurement has a value between x and $x + dx$. We suppose that the PDF of the data has a normal distribution. Most quantitative research involves the use of statistical methods presuming *independence* among data points and Gaussian 'normal' distributions (Andriani and McKelvey, 2007). The Gaussian distribution is reliably characterized by its stable mean and finite variance (Greene, 2002). Normal distributions place a trivial amount of probability far from the mean and hence the mean is representative of most observations. Even the largest deviations, which are exceptionally rare, are still only about a factor of two from the mean in either direction and are well characterized by quoting a simple standard deviation (Clauset, Shalizi, and Newman, 2007). However, apparently rare real life catastrophic events such as major earth quakes, stock market crashes, heavy rainfall events, etc., occur more frequently than indicated by the normal curve, i.e., they exhibit a probability distribution with a *fat tail*. Fat tails indicate a power law pattern and interdependence. The "tails" of a power-law curve — the regions to either side that correspond to large fluctuations — fall off very slowly in comparison with those of the bell curve (Buchanan, 2004). The normal distribution is therefore an inadequate model for extreme departures from the mean.

The following references are cited by Goertzel and Fashing (1981, 1986) to show that the bell curve is an empirical model without supporting theoretical basis: (i) Modern texts usually recognize that there is no theoretical justification for the use of the normal curve, but justify using it as a convenience (Cronbach, 1970). (ii) The bell curve came to be generally accepted, as M. Lippman remarked to Poincare (Bradley, 1969), because "...the experimenters fancy that it is a theorem in mathematics and the mathematicians that it is an experimental fact". (iii) Karl Pearson (best known today for the invention of the product-moment correlation coefficient) used his newly

developed *Chi Square* test to check how closely a number of empirical distributions of supposedly random errors fitted the bell curve. He found that many of the distributions that had been cited in the literature as fitting the normal curve were actually significantly different from it, and concluded that "the normal curve of error possesses no special fitness for describing errors or deviations such as arise either in observing practice or in nature" (Pearson, 1900).

2.2 Fractal fluctuations and statistical analysis

Fractals are the latest development in statistics. The space-time fluctuation pattern in dynamical systems was shown to have a selfsimilar or fractal structure in the 1970s (Mandelbrot, 1975). The larger scale fluctuation consists of smaller scale fluctuations identical in shape to the larger scale. An appreciation of the properties of fractals is changing the most basic ways we analyze and interpret data from experiments and is leading to new insights into understanding physical, chemical, biological, psychological, and social systems. Fractal systems extend over many scales and so cannot be characterized by a single characteristic average number (Liebovitch and Scheurle, 2000). Further, the selfsimilar fluctuations imply long-range space-time correlations or interdependence. Therefore, the Gaussian distribution will not be applicable for description of fractal data sets. However, the bell curve still continues to be used for approximate quantitative characterization of data which are now identified as fractal space-time fluctuations.

2.2.1 Power laws and fat tails

Fractals conform to power laws. A power law is a relationship in which one quantity A is proportional to another B taken to some power n ; that is, $A \sim B^n$ (Buchanan, 2004). One of the oldest scaling laws in geophysics is the Omori law (Omori, 1895). This law describes the temporal distribution of the number of after-shocks, which occur after a larger earthquake (i.e., main-shock) by a scaling relationship. Richardson (1960) came close to the concept of fractals when he noted that the estimated length of an irregular coastline scales with the length of the measuring unit. Andriani and McKelvey (2007) have given exhaustive references to earliest known work on power law relationships summarized as follows. Pareto (1897) first noticed power laws and fat tails in economics. Cities follow a power law when ranked by population (Auerbach, 1913). Dynamics of earthquakes follow power law (Gutenberg and Richter, 1944) and Zipf (1949) found that a power law applies to word frequencies (Estoup (1916), had earlier found a similar relationship). Mandelbrot (1963) rediscovered them in the 20th century, spurring a small wave of interest in finance (Fama, 1965; Montroll and Shlesinger, 1984). However, the rise of the 'standard' model (Gaussian) of efficient markets, sent power law models into obscurity. This lasted until the 1990s, when the occurrence of catastrophic events, such as the 1987 and 1998 financial crashes, that were difficult to explain with the 'standard' models (Bouchaud *et al.*, 1998), re-kindled the fractal model (Mandelbrot and Hudson, 2004).

Sornette (1995) cites the works of Mandelbrot (1983), Aharony and Feder (1989) and Riste and Sherrington (1991) and states that observation that many natural phenomena have size distributions that are power laws, has been taken as a fundamental indication of an underlying self-similarity. A power law distribution indicates the absence of a characteristic size and as a consequence that there is no upper limit on the size of events. The largest events of a power law distribution completely dominate the underlying physical process; for instance, fluid-driven erosion is dominated by the largest floods and most deformation at plate boundaries

takes place through the agency of the largest earthquakes. It is a matter of debate whether power law distributions, which are valid descriptions of the numerous small and intermediate events, can be extrapolated to large events; the largest events are, almost by definition, undersampled.

A power law world is dominated by extreme events ignored in a Gaussian-world. In fact, the fat tails of power law distributions make large extreme events orders-of-magnitude more likely. Theories explaining power laws are also scale-free. This is to say, the same explanation (theory) applies at all levels of analysis (Andriani and McKelvey, 2007).

2.2.2 Scale-free theory for power laws with fat, long tails

A scale-free theory for the observed fractal fluctuations in atmospheric flows shows that the observed long-range spatiotemporal correlations are intrinsic to quantumlike chaos governing fluid flows. The model concepts are independent of the exact details such as the chemical, physical, physiological and other properties of the dynamical system and therefore provide a general systems theory applicable to all real world and computed dynamical systems in nature (Selvam, 1998, 1999, 2001a, b, 2002a, b, 2004, 2005, 2007; Selvam *et al.*, 2000). The model is based on the concept that the irregular fractal fluctuations may be visualized to result from the superimposition of an eddy continuum, i.e., a hierarchy of eddy circulations generated at each level by the space-time integration of enclosed small-scale eddy fluctuations. Such a concept of space-time fluctuation averaged distributions *should* follow statistical normal distribution according to *Central Limit Theorem* in traditional Statistical theory (Ruhla, 1992). Also, traditional statistical/mathematical theory predicts that the Gaussian, its Fourier transform and therefore Fourier transform associated power spectrum are the same distributions. The Fourier transform of normal distribution is essentially a normal distribution. A power spectrum is based on the Fourier transform, which expresses the relationship between time (space) domain and frequency domain description of any physical process (Phillips, 2005; Riley, Hobson and Bence, 2006). However, the model (Sec. 3) visualises the eddy growth process in successive stages of unit length-step growth with ordered two-way energy feedback between the larger and smaller scale eddies and derives a power law probability distribution P which is close to the Gaussian for small deviations and gives the observed fat, long tail for large fluctuations. Further, the model predicts the power spectrum of the eddy continuum also to follow the power law probability distribution P .

In summary, the model predicts the following:

- The eddy continuum consists of an overall logarithmic spiral trajectory with the quasiperiodic *Penrose* tiling pattern for the internal structure.
- The successively larger eddy space-time scales follow the Fibonacci number series.
- The probability distribution P of fractal domains for the n^{th} step of eddy growth is equal to τ^{-4n} where τ is the golden mean equal to $(1+\sqrt{5})/2$ (≈ 1.618). The eddy growth step n represents the *normalized deviation* t in traditional statistical theory. The *normalized deviation* t represents the departure of the variable from the mean in terms of the standard deviation of the distribution assumed to follow *normal distribution* characteristics for many real world space-time events. There is progressive decrease in the probability of occurrence of events with increase in corresponding *normalized deviation* t . Space-time events with *normalized deviation* t greater than 2 occur with a probability of 5 percent or less and may be categorized as extreme events

associated in general with widespread (space-time) damage and loss. The model predicted probability distribution P is close to the statistical normal distribution for t values less than 2 and greater than normal distribution for t more than 2, thereby giving a *fat, long tail*. There is non-zero probability of occurrence of very large events.

- The inverse of probability distribution P , namely, τ^{4n} represents the relative eddy energy flux in the large eddy fractal (small scale fine structure) domain. There is progressive decrease in the probability of occurrence of successive stages of eddy growth associated with progressively larger domains of fractal (small scale fine structure) eddy energy flux and at sufficiently large growth stage trigger catastrophic extreme events such as heavy rainfall, stock market crashes, traffic jams, etc., in real world situations.
- The power spectrum of fractal fluctuations also follows the same distribution P as for the distribution of fractal fluctuations. The square of the eddy amplitude (variance) represents the eddy energy and therefore the eddy probability density P . Such a result that the additive amplitudes of eddies when squared represent probabilities, is exhibited by the sub-atomic dynamics of quantum systems such as the electron or proton (Maddox, 1988, 1993; Rae, 1988). Therefore fractal fluctuations are signatures of quantumlike chaos in dynamical systems.
- The *fine structure constant* for spectrum of fractal fluctuations is a function of the *golden mean* and is analogous to that of atomic spectra equal to about 1/137.
- The universal algorithm for self-organized criticality is expressed in terms of the universal *Feigenbaum's* constants (Feigenbaum, 1980) a and d as $2a^2 = \pi d$ where the fractional volume intermittency of occurrence πd contributes to the total variance $2a^2$ of fractal structures. The *Feigenbaum's* constants are expressed as functions of the *golden mean*. The probability distribution P of fractal domains is also expressed in terms of the *Feigenbaum's* constants a and d . The details of the model are summarized in the following section (Sec. 3).

3. A general systems theory for fractal fluctuations

The fractal space-time fluctuations of dynamical systems may be visualized to result from the superimposition of an ensemble of eddies (sine waves), namely an eddy continuum. The relationship between large and small eddy circulation parameters are obtained on the basis of Townsend's (1956) concept that large eddies are envelopes enclosing turbulent eddy (small-scale) fluctuations (Fig. 2).




Fig. 2: Physical concept of eddy growth process by the self-sustaining process of ordered energy feedback between the larger and smaller scales, the smaller scales forming the internal circulations of larger scales. The figure shows a uniform distribution of dominant turbulent scale eddies of length scale $2r$. Larger-eddy circulations such as ABCD form as coherent structures sustained by the enclosed turbulent eddies.

The relationship between root mean square (r. m. s.) circulation speeds W and w_* respectively of large and turbulent eddies of respective radii R and r is then given as

$$W^2 = \frac{2}{\pi} \frac{r}{R} w_*^2 \quad (1)$$

The dynamical evolution of space-time fractal structures is quantified in terms of ordered energy flow between fluctuations of all scales in Eq. (1), because the square of the eddy circulation speed represents the eddy energy (kinetic). A hierarchical continuum of eddies is generated by the integration of successively larger enclosed turbulent eddy circulations. Such a concept of space-time fluctuation averaged distributions *should* follow statistical normal distribution according to *Central Limit Theorem* in traditional Statistical theory (Ruhla, 1992). Also, traditional statistical/mathematical theory predicts that the Gaussian, its Fourier transform and therefore Fourier transform associated power spectrum are the same distributions. However, the general systems theory (Selvam, 1998, 1999, 2001a, 2001b, 2002a, 2002b, 2004, 2005, 2007; Selvam *et al.*, 2000) visualises the eddy growth process in successive stages of unit length-step growth with ordered two-way energy feedback between the larger and smaller scale eddies and derives a power law probability distribution P which is close to the Gaussian for small deviations and gives the observed fat, long tail for large fluctuations. Further, the model predicts the power spectrum of the eddy continuum also to follow the power law probability distribution P . Therefore the additive amplitudes of the eddies when squared (variance), represent

the probability distribution similar to the subatomic dynamics of quantum systems such as the electron or photon. Fractal fluctuations therefore exhibit quantumlike chaos.

The above-described analogy of quantumlike mechanics for dynamical systems is similar to the concept of a subquantum level of fluctuations whose space-time organization gives rise to the observed manifestation of subatomic phenomena, i.e., quantum systems as order out of chaos phenomena (Grossing, 1989).

3.1 Quasicrystalline structure of the eddy continuum

The turbulent eddy circulation speed and radius increase with the progressive growth of the large eddy (Selvam, 1990, 2007). The successively larger turbulent fluctuations, which form the internal structure of the growing large eddy, may be computed (Eq. 1) as

$$w_*^2 = \frac{\pi R}{2 dR} W^2 \quad (2)$$

During each length step growth dR , the small-scale energizing perturbation W_n at the n^{th} instant generates the large-scale perturbation W_{n+1} of radius R where $R = \sum_1^n dR$ since successive length-scale doubling gives rise to R . Eq. 2 may be written in terms of the successive turbulent circulation speeds W_n and W_{n+1} as

$$W_{n+1}^2 = \frac{\pi R}{2 dR} W_n^2 \quad (3)$$

The angular turning $d\theta$ inherent to eddy circulation for each length step growth is equal to dR/R . The perturbation dR is generated by the small-scale acceleration W_n at any instant n and therefore $dR=W_n$. Starting with the unit value for dR the successive W_n , W_{n+1} , R , and $d\theta$ values are computed from Eq. 3 and are given in Table 1.

Table 1. The computed spatial growth of the strange-attractor design traced by the macro-scale dynamical system of atmospheric flows as shown in Fig. 3.

R	W_n	dR	$d\theta$	W_{n+1}	θ
1.000	1.000	1.000	1.000	1.254	1.000
2.000	1.254	1.254	0.627	1.985	1.627
3.254	1.985	1.985	0.610	3.186	2.237
5.239	3.186	3.186	0.608	5.121	2.845
8.425	5.121	5.121	0.608	8.234	3.453
13.546	8.234	8.234	0.608	13.239	4.061
21.780	13.239	13.239	0.608	21.286	4.669
35.019	21.286	21.286	0.608	34.225	5.277
56.305	34.225	34.225	0.608	55.029	5.885
90.530	55.029	55.029	0.608	88.479	6.493

It is seen that the successive values of the circulation speed W and radius R of the growing turbulent eddy follow the Fibonacci mathematical number series such that $R_{n+1}=R_n+R_{n-1}$ and R_{n+1}/R_n is equal to the golden mean τ , which is equal to $[(1 + \sqrt{5})/2] \cong (1.618)$. Further, the successive W and R values form the geometrical progression $R_0(1+\tau+\tau^2+\tau^3+\tau^4+ \dots)$ where R_0 is the initial value of the turbulent eddy radius.

Turbulent eddy growth from primary perturbation OR_0 starting from the origin O (Fig. 3) gives rise to compensating return circulations OR_1R_2 on either side of OR_0 ,

thereby generating the large eddy radius OR_1 such that $OR_1/OR_0=\tau$ and $R_0OR_1=\pi/5=R_0R_1O$. Therefore, short-range circulation balance requirements generate successively larger circulation patterns with precise geometry that is governed by the *Fibonacci* mathematical number series, which is identified as a signature of the universal period doubling route to chaos in fluid flows, in particular atmospheric flows. It is seen from Fig. 3 that five such successive length step growths give successively increasing radii OR_1 , OR_2 , OR_3 , OR_4 and OR_5 tracing out one complete vortex-roll circulation such that the scale ratio OR_5/OR_0 is equal to $\tau^5=11.1$. The envelope $R_1R_2R_3R_4R_5$ (Fig. 3) of a dominant large eddy (or vortex roll) is found to fit the logarithmic spiral $R=R_0e^{b\theta}$ where $R_0=OR_0$, $b=\tan \delta$ with δ the crossing angle equal to $\pi/5$, and the angular turning θ for each length step growth is equal to $\pi/5$. The successively larger eddy radii may be subdivided again in the *golden mean* ratio. The internal structure of large-eddy circulations is therefore made up of balanced small-scale circulations tracing out the well-known quasi-periodic *Penrose* tiling pattern identified as the quasi-crystalline structure in condensed matter physics. A complete description of the atmospheric flow field is given by the quasi-periodic cycles with *Fibonacci* winding numbers.

3.2 Model predictions

The model predictions (Selvam, 1990, 2005, 2007; Selvam and Fadnavis, 1998) are

3.2.1 Quasiperiodic Penrose tiling pattern

Atmospheric flows trace an overall logarithmic spiral trajectory $OR_0R_1R_2R_3R_4R_5$ simultaneously in clockwise and anti-clockwise directions with the quasi-periodic *Penrose* tiling pattern (Steinhardt, 1997) for the internal structure shown in Fig. 3.




Fig. 3. The quasiperiodic *Penrose* tiling pattern

(clockwise and anti-clockwise) directions. Eddy fluctuation length of amplitude $1/\tau$ has a probability of occurrence equal to $1/\tau^2$ in both rotational directions, i.e., the square of eddy amplitude represents the probability of occurrence in the eddy continuum. Similar result is observed in the subatomic dynamics of quantum systems which are visualized to consist of the superimposition of eddy fluctuations in wave trains (eddy continuum).

The spiral flow structure can be visualized as an eddy continuum generated by successive length step growths OR_0 , OR_1 , OR_2 , OR_3 ,... respectively equal to R_1 , R_2 , R_3 ,... which follow *Fibonacci* mathematical series such that $R_{n+1}=R_n+R_{n-1}$ and $R_{n+1}/R_n=\tau$ where τ is the *golden mean* equal to $(1+\sqrt{5})/2$ (≈ 1.618). Considering a normalized length step equal to 1 for the last stage of eddy growth, the successively decreasing radial length steps can be expressed as 1 , $1/\tau$, $1/\tau^2$, $1/\tau^3$, The normalized eddy continuum comprises of fluctuation length scales 1 , $1/\tau$, $1/\tau^2$, The probability of occurrence is equal to $1/\tau$ and $1/\tau^2$ respectively for eddy length scale $1/\tau$ in any one or both rotational

3.2.2 Logarithmic spiral pattern underlying fractal fluctuations

The overall logarithmic spiral flow structure (Fig. 1) is given by the relation

$$W = \frac{w_*}{k} \ln z \quad (4)$$

In Eq. (4) the constant k is the steady state fractional volume dilution of large eddy by inherent turbulent eddy fluctuations and z is the length scale ratio R/r . The constant k is equal to $1/\tau^2$ ($\cong 0.382$) and is identified as the universal constant for deterministic chaos in fluid flows. The steady state emergence of fractal structures is therefore equal to

$$\frac{1}{k} \cong 2.62 \quad (5)$$

In Eq. (4), W represents the standard deviation of eddy fluctuations, since W is computed as the instantaneous r. m. s. (root mean square) eddy perturbation amplitude with reference to the earlier step of eddy growth. For two successive stages of eddy growth starting from primary perturbation w_* , the ratio of the standard deviations W_{n+1} and W_n is given from Eq. (4) as $(n+1)/n$. Denoting by σ the standard deviation of eddy fluctuations at the reference level ($n=1$) the standard deviations of eddy fluctuations for successive stages of eddy growth are given as integer multiples of σ , i.e., σ , 2σ , 3σ , etc. and correspond respectively to

$$\text{statistical normalised standard deviation } t = 0, 1, 2, 3, \dots \quad (6)$$

The conventional power spectrum plotted as the variance versus the frequency in log-log scale will now represent the eddy probability density on logarithmic scale versus the standard deviation of the eddy fluctuations on linear scale since the logarithm of the eddy wavelength represents the standard deviation, i.e., the r. m. s. value of eddy fluctuations (Eq. 4). The r. m. s. value of eddy fluctuations can be represented in terms of statistical normal distribution as follows. A normalized standard deviation $t=0$ corresponds to cumulative percentage probability density equal to 50 for the mean value of the distribution. Since the logarithm of the wavelength represents the r. m. s. value of eddy fluctuations the normalized standard deviation t is defined for the eddy energy as

$$t = \frac{\log L}{\log T_{50}} - 1 \quad (7)$$

In Eq. (7) L is the time period (or wavelength) and T_{50} is the period up to which the cumulative percentage contribution to total variance is equal to 50 and $t = 0$. $\log T_{50}$ also represents the mean value for the r. m. s. eddy fluctuations and is consistent with the concept of the mean level represented by r. m. s. eddy fluctuations. Spectra of time series of meteorological parameters when plotted as cumulative percentage contribution to total variance versus t have been shown to follow closely the model predicted universal spectrum (Selvam and Fadnavis, 1998) which is identified as a signature of quantumlike chaos.

3.2.3 Universal Feigenbaum's constants and probability distribution function for fractal fluctuations

Selvam (1993, 2007) has shown that Eq. (1) represents the universal algorithm for deterministic chaos in dynamical systems and is expressed in terms of the universal Feigenbaum's (1980) constants a and d as follows. The successive length step growths generating the eddy continuum $OR_0R_1R_2R_3R_4R_5$ (Fig. 1) analogous to the period doubling route to chaos (growth) is initiated and sustained by the turbulent (fine scale) eddy acceleration w_* , which then propagates by the inherent property of inertia of the medium of propagation. Therefore, the statistical parameters *mean*, *variance*, *skewness* and *kurtosis* of the perturbation field in the medium of propagation are given by w_* , w_*^2 , w_*^3 and w_*^4 respectively. The associated dynamics of the perturbation field can be described by the following parameters. The perturbation speed w_* (motion) per second (unit time) sustained by its inertia represents the mass, w_*^2 the acceleration or force, w_*^3 the angular momentum or potential energy, and w_*^4 the spin angular momentum, since an eddy motion has an inherent curvature to its trajectory.

It is shown that Feigenbaum's constant a is equal to (Selvam, 1993, 2007)

$$a = \frac{W_2R_2}{W_1R_1} \quad (8)$$

In Eq. (8) the subscripts 1 and 2 refer to two successive stages of eddy growth. Feigenbaum's constant a as defined above represents the steady state emergence of fractional *Euclidean* structures. Considering dynamical eddy growth processes, Feigenbaum's constant a also represents the steady state fractional outward mass dispersion rate and a^2 represents the energy flux into the environment generated by the persistent primary perturbation W_1 . Considering both clockwise and counterclockwise rotations, the total energy flux into the environment is equal to $2a^2$. In statistical terminology, $2a^2$ represents the variance of fractal structures for both clockwise and counterclockwise rotation directions.

The probability of occurrence P_{tot} of fractal domain W_1R_1 in the total larger eddy domain W_nR_n in any (irrespective of positive or negative) direction is equal to

$$P_{tot} = \frac{W_1R_1}{W_nR_n} = \tau^{-2n}$$

Therefore the probability P of occurrence of fractal domain W_1R_1 in the total larger eddy domain W_nR_n in any one direction (either positive or negative) is equal to

$$P = \left(\frac{W_1R_1}{W_nR_n} \right)^2 = \tau^{-4n} \quad (9)$$

The Feigenbaum's constant d is shown to be equal to (Selvam, 1993, 2007)

$$d = \frac{W_2^4R_2^3}{W_1^4R_1^3} \quad (10)$$

Eq. (10) represents the fractional volume intermittency of occurrence of fractal structures for each length step growth. *Feigenbaum's* constant d also represents the relative spin angular momentum of the growing large eddy structures as explained earlier.

Eq. (1) may now be written as

$$2 \frac{W^2 R^2}{w_*^2 (dR)^2} = \pi \frac{W^4 R^3}{w_*^4 (dR)^3} \quad (11)$$

In Eq. (11) dR equal to r represents the incremental growth in radius for each length step growth, i.e., r relates to the earlier stage of eddy growth.

The Feigenbaum's constant d represented by R/r is equal to

$$d = \frac{W^4 R^3}{w_*^4 r^3} \quad (12)$$

For two successive stages of eddy growth

$$d = \frac{W_2^4 R_2^3}{W_1^4 R_1^3} \quad (13)$$

From Eq. (1)

$$W_1^2 = \frac{2}{\pi} \frac{r}{R_1} w_*^2 \quad (14)$$

$$W_2^2 = \frac{2}{\pi} \frac{r}{R_2} w_*^2$$

Therefore

$$\frac{W_2^2}{W_1^2} = \frac{R_1}{R_2} \quad (15)$$

Substituting in Eq. (13)

$$d = \frac{W_2^4 R_2^3}{W_1^4 R_1^3} = \frac{W_2^2}{W_1^2} \frac{W_2^2 R_2^3}{W_1^2 R_1^3} = \frac{R_1}{R_2} \frac{W_2^2 R_2^3}{W_1^2 R_1^3} = \frac{W_2^2 R_2^2}{W_1^2 R_1^2} \quad (16)$$

The Feigenbaum's constant d represents the scale ratio R_2/R_1 and the inverse of the Feigenbaum's constant d equal to R_1/R_2 represents the probability $(Prob)_1$ of occurrence of length scale R_1 in the total fluctuation length domain R_2 for the first eddy growth step as given in the following

$$(Prob)_1 = \frac{R_1}{R_2} = \frac{1}{d} = \frac{W_1^2 R_1^2}{W_2^2 R_2^2} = \tau^{-4} \quad (17)$$

In general for the n^{th} eddy growth step, the probability $(Prob)_n$ of occurrence of length scale R_1 in the total fluctuation length domain R_n is given as

$$(Prob)_n = \frac{R_1}{R_n} = \frac{W_1^2 R_1^2}{W_n^2 R_n^2} = \tau^{-4n} \quad (18)$$

The above equation for probability $(Prob)_n$ also represents, for the n^{th} eddy growth step, the following statistical and dynamical quantities of the growing large eddy with respect to the initial perturbation domain: (i) the statistical relative variance of fractal structures, (ii) probability of occurrence of fractal domain in either positive or negative direction, and (iii) the inverse of $(Prob)_n$ represents the organised fractal (fine scale) energy flux in the overall large scale eddy domain. Large scale energy flux therefore occurs not in bulk, but in organized internal fine scale circulation structures identified as fractals.

Substituting the *Feigenbaum's constants* a and d defined above (Eqs. 8 and 10), Eq. (11) can be written as

$$2a^2 = \pi d \quad (19)$$

In Eq. (19) πd , the relative volume intermittency of occurrence contributes to the total variance $2a^2$ of fractal structures.

In terms of eddy dynamics, the above equation states that during each length step growth, the energy flux into the environment equal to $2a^2$ contributes to generate relative spin angular momentum equal to πd of the growing fractal structures. Each length step growth is therefore associated with a factor of $2a^2$ equal to $2\tau^4$ ($\cong 13.70820393$) increase in energy flux in the associated fractal domain. Ten such length step growths results in the formation of robust (self-sustaining) dominant bidirectional large eddy circulation $OR_0R_1R_2R_3R_4R_5$ (Fig. 1) associated with a factor of $20a^2$ equal to 137.08203 increase in eddy energy flux. This non-dimensional constant factor characterizing successive dominant eddy energy increments is analogous to the *fine structure* constant α^{-1} (Ford, 1968) observed in atomic spectra, where the spacing (energy) intervals between adjacent spectral lines is proportional to the non-dimensional *fine structure* constant equal to approximately 1/137. Further, the probability of n^{th} length step eddy growth is given by a^{-2n} ($\cong 6.8541^{-n}$) while the associated increase in eddy energy flux into the environment is equal to a^{2n} ($\cong 6.8541^n$). Extreme events occur for large number of length step growths n with small probability of occurrence and are associated with large energy release in the fractal domain. Each length step growth is associated with one-tenth of *fine structure constant* energy increment equal to $2a^2$ ($\alpha^{-1}/10 \cong 13.7082$) for bidirectional eddy circulation, or equal to one-twentieth of *fine structure constant* energy increment equal to a^2 ($\alpha^{-1}/20 \cong 6.8541$) in any one direction, i.e., positive or negative. The energy increase between two successive eddy length step growths may be expressed as a function of $(a^2)^2$, i.e., proportional to the square of the *fine structure constant* α^{-1} . In the spectra of many atoms, what appears with coarse observations to be a single spectral line proves, with finer observation, to be a group of two or more closely spaced lines. The spacing of these fine-structure lines relative to the coarse spacing in the spectrum is proportional to the square of *fine structure constant*, for which reason this combination is called the *fine-structure constant*. We now know that the significance of the *fine-structure constant* goes beyond atomic spectra (Ford, 1968).

It was shown at Eq. (5) (Sec. 3.2.2) above that the steady state emergence of fractal structures in fluid flows is equal to $1/k (= \tau^2)$ and therefore the *Feigenbaum's constant* a is equal to

$$a = \tau^2 = \frac{1}{k} = 2.62 \quad (20)$$

3.2.4 Universal Feigenbaum's constants and power spectra of fractal fluctuations

The power spectra of fluctuations in fluid flows can now be quantified in terms of universal *Feigenbaum's constant* a as follows.

The normalized variance and therefore the statistical probability distribution is represented by (from Eq. 9)

$$P = a^{-2t} \quad (21)$$

In Eq. (21) P is the probability density corresponding to normalized standard deviation t . The graph of P versus t will represent the power spectrum. The slope S of the power spectrum is equal to

$$S = \frac{dP}{dt} \approx -P \quad (22)$$

The power spectrum therefore follows inverse power law form, the slope decreasing with increase in t . Increase in t corresponds to large eddies (low frequencies) and is consistent with observed decrease in slope at low frequencies in dynamical systems.

The probability distribution of fractal fluctuations (Eq. 18) is therefore the same as variance spectrum (Eq. 21) of fractal fluctuations.

The steady state emergence of fractal structures for each length step growth for any one direction of rotation (either clockwise or anticlockwise) is equal to

$$\frac{a}{2} = \frac{\tau^2}{2}$$

since the corresponding value for both direction is equal to a (Eqs. 5 and 20).

The emerging fractal space-time structures have moment coefficient of kurtosis given by the fourth moment equal to

$$\left(\frac{\tau^2}{2} \right)^4 = \frac{\tau^8}{16} = 2.9356 \approx 3$$

The moment coefficient of skewness for the fractal space-time structures is equal to zero for the symmetric eddy circulations. Moment coefficient of kurtosis equal to 3 and moment coefficient of skewness equal to *zero* characterize the statistical normal distribution. The model predicted power law distribution for fractal fluctuations is close to the Gaussian distribution.

3.2.5 The power spectrum and probability distribution of fractal fluctuations are the same

The relationship between *Feigenbaum's constant* a and power spectra may also be derived as follows.

The steady state emergence of fractal structures is equal to the *Feigenbaum's constant* a (Eqs. 5 and 20). The relative variance of fractal structure which also represents the probability P of occurrence of bidirectional fractal domain for each length step growth is then equal to $1/a^2$. The normalized variance $\frac{1}{a^{2n}}$ will now represent the statistical probability density for the n^{th} step growth according to model predicted quantumlike mechanics for fluid flows. Model predicted probability density values P are computed as

$$P = \frac{1}{a^{2n}} = \tau^{-4n} \quad (23)$$

or

$$P = \tau^{-4t} \quad (24)$$

In Eq. (24) t is the normalized standard deviation (Eq. 6). The model predicted P values corresponding to normalised deviation t values less than 2 are slightly less than the corresponding statistical normal distribution values while the P values are noticeably larger for normalised deviation t values greater than 2 (Table 2 and Fig. 4) and may explain the reported *fat tail* for probability distributions of various physical parameters (Buchanan, 2004). The model predicted P values plotted on a linear scale (Y-axis) shows close agreement with the corresponding statistical normal probability values as seen in Fig.4 (left side). The model predicted P values plotted on a logarithmic scale (Y-axis) shows *fat tail* distribution for normalised deviation t values greater than 2 as seen in Fig.4 (right side).

Table 2: Model predicted and statistical normal probability density distributions

growth step	normalized deviation	cumulative probability densities (%)	
n	t	model predicted $P = \tau^{-4t}$	statistical normal distribution
1	1	14.5898	15.8655
2	2	2.1286	2.2750
3	3	0.3106	0.1350
4	4	0.0453	0.0032
5	5	0.0066	≈ 0.0

comparison of statistical normal distribution and computed (theoretical) probability density distribution




Fig. 4: Comparison of statistical normal distribution and computed (theoretical) probability density distribution. The same figure is plotted on the right side with logarithmic scale for the probability axis (Y-axis) to show clearly that for normalized deviation t values greater than 2 the computed probability densities are greater than the corresponding statistical normal distribution values.

4. Data and analysis

4.1 Data sets used for the study

Gridded temperature anomalies for mean temperatures obtained from the GHCN V2 (<http://www.ncdc.noaa.gov/oa/climate/ghcn-monthly/index.php> and <ftp://ftp.ncdc.noaa.gov/pub/data/ghcn/v2/>) monthly temperature data sets for 129 years (1880 to 2008) the months January to May and for 128 years (1880 to 2008) for the months June to December were used for the study. Details of the data set as given in the 'README_GRID_TEMP' text document are as follows. GHCN homogeneity adjusted data was the primary source for developing the gridded fields. In grid boxes without homogeneity adjusted data, GHCN raw data was used to provide additional coverage when possible. Each month of data consists of 2592 gridded data points produced on a 5 X 5 degree basis for the entire globe (72 longitude X 36 latitude grid boxes).

Gridded data for every month from January 1880 to the most recent month (May 2008 in the present study) is available. The data are temperature anomalies in degrees Celsius. Each gridded value was multiplied by 100 and written to file as an integer. Missing values are represented by the value -32768.

There are two options: `grid_1880_YYYY.dat.gz` (used in the present study), where YYYY is the current year and the values are calculated using the first difference method, and `anom-grid-1880-current.dat.gz` which uses the "anomaly" method.

The data are formatted by year, month, latitude and longitude. There are twelve longitude grid values per line, so there are 6 lines ($72/12 = 6$) for each of the 36 latitude bands. Longitude values are written from 180 W to 180 E, and latitude values

from 90 N to 90 S. Data for each month is preceded by a label containing the month and year of the gridded data.

```

for year = begyr to endyr
  for month = 1 to 12
    format(2i5) month,year
    for ylat = 1 to 36 (85-90N,80-85N,....,80-85S,85-90S)
      format(12i5) 180-175W,175-170W,....,130-125W,125-120W
      format(12i5) 120-115W,175-170W,....,70-65W,65-60W
      format(12i5) 60-55W,55-50W,....,10-5W,5-0W
      format(12i5) 0-5E,5-10E,....,50-55E,55-60E
      format(12i5) 60-65E,65-70E,....,110-115E,115-120E
      format(12i5) 120-125E,125-130E,....,170-175E,175-180E

```

Each file has been compressed using 'gzip'. They can be uncompressed with 'WinZip' for those using Windows 95 (and above) or with 'gzip' from most UNIX platforms. The FORTRAN utility program 'read_gridded.f' can be downloaded to assist in extracting data of interest. This program allows the user to extract non-missing values for selected months and write the data to an ascii output file. The latitude and longitude of the center of each corresponding grid box accompanies each gridded value in the output file.

We calculated these anomalies with respect to the period 1961 - 1990 using the First Difference Method, an approach developed to maximize the use of available station records (Peterson and Vose, 1997; Peterson et al., 1998a; Peterson et al., 1998b). The First Difference Method involves calculating a series of calendar-month differences in temperature between successive years of station data ($FDyr = Tyr - Tyr-1$). For example, when creating a station's first difference series for mean February temperature, we subtract the station's February 1880 temperature from the station's February 1881 temperature to create a February 1881 first difference value. First difference values for subsequent years are calculated in the same fashion by subtracting the station's preceding year temperature for all available years of station data.

For each year and month we sum the 'first difference' value of all stations located within the appropriate 5 X 5 degree box and divide by the total number of stations in the box to get an unweighted first difference value for each grid box. We then calculate a cumulative sum of these gridded first difference values for all years from 1880 to 1998 to produce a time series for each grid box. The cumulative sum is calculated for each grid box and each month of gridded first difference data independently through time. Each grid box time series is then adjusted to create anomalies with respect to the period 1961 - 1990.

We developed this gridded data set to produce the most accurate time series possible. However, this required that we treat months and grid boxes independently through time. The use of this data is most appropriate for analyzing the change in temperature within a particular grid box, or set of grid boxes, over a span of years. If one is more interested in analyzing temperature changes within individual years, e.g., the change in temperature between February and March, 1908, or between two regions in 1908, we recommend that the GHCN station data be used directly.

4.2 Analyses and results

4.2.1 Frequency distribution

Each data set (month-wise temperature time series for the period 1800 to 2008 for the months January to May and for the period 1800 to 2007 for the months June to December) was represented as the frequency of occurrence $f(i)$ in a suitable number n of class intervals $x(i)$, $i=1, n$ covering the range of values from *minimum* to the *maximum* in the data set. The class interval $x(i)$ represents dataset values in the range $x(i) \pm \Delta x$, where Δx is a constant. The average av and standard deviation sd for the data set is computed as

$$av = \frac{\sum_1^n [x(i) \times f(i)]}{\sum_1^n f(i)}$$

$$sd = \frac{\sum_1^n \{ [x(i) - av]^2 \times f(i) \}}{\sum_1^n f(i)}$$

The *normalized deviation* t values for class intervals $t(i)$ were then computed as

$$t(i) = \frac{x(i) - av}{sd}$$

The cumulative percentage probabilities of occurrence $cmax(i)$ and $cmin(i)$ corresponding to the *normalized deviation* t values were then computed starting respectively from the maximum ($i=n$) and minimum ($i=1$) class interval values as follows.

$$cmax(i) = \frac{\sum_1^i [x(i) \times f(i)]}{\sum_1^n [x(i) \times f(i)]} \times 100.0$$

$$cmin(i) = \frac{\sum_1^i [x(i) \times f(i)]}{\sum_1^n [x(i) \times f(i)]} \times 100.0$$

The $cmax$ and $cmin$ distributions with respect to the *normalized deviation* t values were computed for each month for all grid points including grid points which did not have continuous time series data. The total number of grid points available for the study for the months January to December is given in Fig. 5. The average and standard deviation of $cmax(i)$ and $cmin(i)$ for each *normalized deviation* $t(i)$ values were then computed for each month from the corresponding distributions for all the available grid points for the month. The average cumulative percentage probability values $cmax(i)$ and $cmin(i)$ plotted with respect to corresponding *normalized deviation* $t(i)$ values with linear axes are shown in Fig. 6 and on logarithmic scale for the

probability axis in the tail region, i.e. *normalized deviation* $t(i)$ values greater than 2 in Fig. 7 for positive extremes $t(i) = 2$ to 4, and in Fig. 8 for negative extremes $t(i) = -2$ to -4 . The standard deviation of each mean $cmax(i)$ and $cmin(i)$ value is shown as a vertical error bar on either side of the mean in Figs. 6 to 8. The figures also contain the statistical normal distribution and the computed theoretical distribution (Eq. 24) for comparison. The figures show clearly the appreciable positive departure of observed probability densities from the statistical normal distribution for extreme values at normalised deviation t values more than 2 (Figs. 7 and 8). The observed extreme values corresponding to $t(i)$ values greater than 2 for $cmax(i)$ and $cmin(i)$ distributions were compared for ‘goodness of fit’ with computed theoretical distribution and statistical normal distribution as follows. For $cmax(i)$ and $cmin(i)$ values, where standard deviation is available (no of observed values more than one), if the observed distribution included the computed theoretical (statistical normal) distribution within twice the standard deviation on either side of the mean then it was assumed to be the same as the computed theoretical (statistical normal) distribution at 5% level of significance within measurement errors. The number of observed distribution values which included the computed theoretical values and/or the statistical normal distribution values within twice the standard deviation on either side of the mean was determined. The total and percentage numbers of observed extreme values same as computed theoretical and statistical normal distributions are given in Fig. 9 for positive and negative tail regions (normalized deviation t greater than 2) for the 12 months January to December. The percentage number of observed extreme value points same as model predicted computed is more than the percentage number of extreme value points same as statistical normal distribution for all the 12 months. The observed distribution is closer to the model predicted theoretical than the statistical normal distribution.

4.2.2 *Continuous periodogram power spectral analyses*

The power spectra of frequency distribution of monthly mean data sets were computed accurately by an elementary, but very powerful method of analysis developed by Jenkinson (1977) which provides a quasi-continuous form of the classical periodogram allowing systematic allocation of the total variance and degrees of freedom of the data series to logarithmically spaced elements of the frequency range $(0.5, 0)$. The cumulative percentage contribution to total variance was computed starting from the high frequency side of the spectrum. The power spectra were plotted as cumulative percentage contribution to total variance versus the normalized standard deviation t equal to $(\log L / \log T_{50}) - 1$ where L is the period in years and T_{50} is the period up to which the cumulative percentage contribution to total variance is equal to 50 (Eq. 7). The statistical *Chi-Square* test (Spiegel, 1961) was applied to determine the ‘goodness of fit’ of variance spectra with statistical normal distribution which is close to model predicted variance spectrum (Eqs. 21 and 24).

gridded temperature anomalies
 Global Historical Climatology Network
 (GHCN-Monthly) data Jan - Dec 1880-2007/2008





Fig. 5: The total number of grid points available for the study for the months January to December.


gridded temperature anomalies January - December 1880-2007/2008
 Global Historical Climatology Network (GHCN-Monthly) data
 mean frequency distribution



- gridded temperature anomalies Jan - Dec 1880-2007/2008 mean frequency distribution error bars (vertical) indicate one standard deviation
- statistical normal distribution — computed (theoretical) distribution

Fig 6: Average cumulative percentage probability distribution


gridded temperature anomalies January - December 1880-2007/2008
Global Historical Climatology Network (GHCN-Monthly) data
mean frequency distribution



- gridded temperature anomalies Jan - Dec 1880-2007/2008 mean frequency distribution error bars (vertical) indicate one standard deviation
- statistical normal distribution ——— computed (theoretical) distribution

Fig 7: Average cumulative percentage probability values on logarithmic scale for the probability axis in the positive tail region (extreme events), i.e., *normalized deviation t* values greater than 2

gridded temperature anomalies January - December 1880-2007/2008
Global Historical Climatology Network (GHCN-Monthly) data
mean frequency distribution



- gridded temperature anomalies Jan - Dec 1880-2007/2008 mean frequency distribution error bars (vertical) indicate one standard deviation
- statistical normal distribution ——— computed (theoretical) distribution

Fig 8: Average cumulative percentage probability values on logarithmic scale for the probability axis in the negative tail region (extreme events), i.e. *normalized deviation t* values greater than -2




Fig 9: The total and percentage numbers of observed extreme values same as model predicted theoretical and statistical normal distributions for positive and negative tail regions (*normalized deviation t* greater than 2) for the 12 months January to December




Fig. 10: Average power (variance) spectrum with vertical error bars indicating one standard deviation on either side of the mean. The statistical normal distribution is also shown in the figure for comparison.




Fig. 11: The total number of grid points and the percentage number of grid points with variance spectra same as statistical normal distribution.

5. Discussion and conclusions

Dynamical systems in nature exhibit selfsimilar fractal fluctuations for all space-time scales and the corresponding power spectra follow inverse power law form signifying long-range space-time correlations identified as self-organized criticality. The physics of self-organized criticality is not yet identified. The Gaussian probability distribution used widely for analysis and description of large data sets is found to significantly underestimate the probabilities of occurrence of extreme events such as stock market crashes, earthquakes, heavy rainfall, etc. Further, the assumptions underlying the normal distribution such as fixed mean and standard deviation, independence of data, are not valid for real world fractal data sets exhibiting a scale-free power law distribution with fat tails. It is important to identify and quantify the fractal distribution characteristics of dynamical systems for predictability studies.

A recently developed general systems theory for fractal space-time fluctuations (Selvam, 1990, 2005, 2007; Selvam and Fadnavis, 1998) shows that the larger scale fluctuation can be visualized to emerge from the space-time averaging of enclosed small scale fluctuations, thereby generating a hierarchy of selfsimilar fluctuations manifested as the observed eddy continuum in power spectral analyses of fractal fluctuations.

The model predictions are as follows.

- The probability distribution function P of fractal fluctuations follow inverse power law form τ^{-4t} where τ is the *golden mean*, and t , the normalized

deviation is equal to $(x-av/sd)$ where av and sd are respectively the average and standard deviation of the distribution. The predicted distribution is close to the Gaussian distribution for small-scale fluctuations (normalized deviation t less than 2), but exhibits *fat long tail* for large-scale fluctuations (normalized deviation t more than 2) with higher probability of occurrence than predicted by Gaussian distribution. There is always a non-zero probability of occurrence of very large amplitude, damage causing extreme events.

- The inverse of the probability distribution function, i.e. $1/P$ equal to τ^{4t} represents the domain size extent of the internal fine scale (fractal) fluctuations of amplitude t (normalized deviation). High intensity extreme events corresponding to t values more than 2 occur with less probability over a larger size domain and are associated with widespread damage and loss such as in heavy rainfall, earthquakes, traffic jams, etc.
- The power spectra of fractal fluctuations (Sec. 3) also follow inverse power law form τ^{-4t} and t , the normalized deviation is expressed in terms of component periodicities L as $t = \{(\log L/\log T_{50})-1\}$ where T_{50} is the period upto which the cumulative percentage contribution to total variance is equal to 50. Since the power (variance, i.e., square of eddy amplitude) spectrum also represents the probability densities as in the case of quantum systems such as the electron or photon, fractal fluctuations exhibit quantumlike chaos.
- The precise geometry of the quasiperiodic Penrose tiling pattern underlie fractal fluctuations tracing out robust (self-sustaining) dominant bidirectional large eddy circulation $OR_0R_1R_2R_3R_4R_5$ (Fig. 1) associated with a factor of $20a^2$ ($20\tau^4$) equal to 137.08203 increase in eddy energy flux. This non-dimensional dominant eddy energy flux is analogous to and almost equal to the *fine structure constant* α^{-1} of atomic spectra; the energy spacing intervals between successive atomic spectral lines. Further, the energy increase between two successive fine structure eddy length step growths internal to the dominant large eddy domain may be expressed as a function of $(a^2)^2$, i.e., proportional to the square of the *fine structure constant* α^{-1} . In the spectra of many atoms, what appears with coarse observations to be a single spectral line proves, with finer observation, to be a group of two or more closely spaced lines. The spacing of these fine-structure lines relative to the coarse spacing in the spectrum is proportional to the square of *fine structure constant*, for which reason this combination is called the *fine-structure constant* (Ford, 1968).

Analysis of historic (1880 -2008) data sets of global monthly mean temperature (GHCN V2) time series show that the data follow closely, but not exactly the statistical normal distribution in the region of normalized deviations t less than 2 (Fig. 6). For normalized deviations t greater than 2, the data exhibit significantly larger probabilities as compared to the normal distribution and closer to the model predicted probability distribution (Figs 7 and 8). A simple t test for ‘goodness of fit’ of the extreme values (normalized deviation $t > 2$) of the observed distribution with model predicted (theoretical) and also the statistical normal distribution shows that more number of data points exhibit significant (at 5% level) ‘goodness of fit’ with the model predicted (theoretical) distribution than with the normal distribution (Fig. 9). The mean power spectra follow closely the statistical normal distribution for the twelve months (Fig. 10). The power spectra mostly cover the region for *normalized deviation* t less than 2 where the model predicted theoretical distribution is close to the statistical normal distribution. A majority (more than 90%) of the power spectra follow closely statistical normal distribution (Fig. 11) consistent with model

prediction of quantumlike chaos, i.e., variance or square of eddy amplitude represents the probability distribution, a signature of quantum systems. The model predicted and observed universal spectrum for interannual variability rules out linear secular trends in global monthly mean temperatures. Global warming results in intensification of fluctuations of all scales and manifested immediately in high frequency fluctuations. The general systems theory, originally developed for turbulent fluid flows, provides universal quantification of physics underlying fractal fluctuations and is applicable to all dynamical systems in nature independent of its physical, chemical, electrical, or any other intrinsic characteristic.

Acknowledgement

The author is grateful to Dr. A. S. R. Murty for encouragement during the course of this study.

References

- A. Aharony and J. Feder, eds, *Fractals in Physics*, (North Holland, Amsterdam), *Physica D* **38** (1989) 1-3.
- P. Andriani and B. McKelvey, Beyond Gaussian averages: Redirecting management research toward extreme events and power laws, *Journal of International Business Studies* **38** (2007) 1212–1230.
- F. Auerbach, Das Gesetz Der Bevölkerungskonzentration, *Petermanns Geographische Mitteilungen* **59** (1913) 74–76.
- P. C. Bak, C. Tang and K. Wiesenfeld, Self-organized criticality, *Phys. Rev. A.* **38** (1988) 364 - 374.
- J. P. Bouchaud, D. Sornette, C. Walter and J. P. Aguilar, Taming large events: Optimal portfolio theory for strongly fluctuating assets, *International Journal of Theoretical and Applied Finance* **1**(1) (1998) 25–41.
- J. V. Bradley, *Distribution-free Statistical Tests*, (Englewood Cliffs, Prentice-Hall, N.J., 1968).
- M. Buchanan, Power laws and the new science of complexity management, *Strategy and Business Issue* **34** (2004) 70-79.
- A. Clauset, C. R. Shalizi and M. E. J. Newman, Power-law distributions in empirical data, arXiv:0706.1062v1 [physics.data-an] 2007.
- L. Cronbach, *Essentials of Psychological Testing*, (Harper & Row, New York, 1970).
- J. B. Estoup, *Gammes Stenographiques*, (Institut Stenographique de France, Paris, 1916).
- E. F. Fama, The behavior of stock-market prices, *Journal of Business* **38** (1965) 34–105.
- M. J. Feigenbaum, Universal behavior in nonlinear systems, *Los Alamos Sci.* **1** (1980) 4-27.
- K. W. Ford, *Basic Physics*, (Blaisdell Publishing Company, Waltham, Massachusetts, USA, 1968, p. 775)
- T. Goertzel and J. Fashing, The myth of the normal curve: A theoretical critique and examination of its role in teaching and research, *Humanity and Society* **5** (1981) 14-31; reprinted in *Readings in Humanist Sociology*, General Hall, 1986. <http://crab.rutgers.edu/~goertzel/normalcurve.htm> 4/29/2007
- W. H. Greene, *Econometric Analysis*, 5th ed. (Prentice-Hall, Englewood Cliffs, NJ, 2002).
- G. Grossing, Quantum systems as order out of chaos phenomena, *Il Nuovo Cimento* **103B** (1989) 497-510.
- B. Gutenberg and R. F. Richter, Frequency of earthquakes in California, *Bulletin of the Seismological Society of America* **34** (1944) 185–188.
- A. F. Jenkinson, *A Powerful Elementary Method of Spectral Analysis for use with Monthly, Seasonal or Annual Meteorological Time Series*, (Meteorological Office, Branch Memorandum No. 57, London, 1977).
- L. S. Liebovitch and D. Scheurle, Two lessons from fractals and chaos, *Complexity* **5**(4) (2000) 34–43.
- J. Maddox, Licence to slang Copenhagen? *Nature* **332** (1988) 581.
- J. Maddox, Can quantum theory be understood? *Nature* **361** (1993) 493.
- B. B. Mandelbrot, The variation of certain speculative prices, *Journal of Business* **36** (1963) 394–419.
- B. B. Mandelbrot, *Les Objets Fractals: Forme, Hasard et Dimension*, (Flammarion, Paris, 1975).
- B. B. Mandelbrot, *The Fractal Geometry of Nature*, (Freeman, New York, 1983).
- B. B. Mandelbrot and R. L. Hudson, *The (Mis)Behavior of Markets: A Fractal View of Risk, Ruin and Reward*, (Profile, London, 2004).
- E. Montroll and M. Shlesinger, On the wonderful world of random walks, in J. L. Lebowitz and E. W. Montroll (eds.) *Nonequilibrium Phenomena II, from Stochastic to Hydrodynamics*, (North Holland, Amsterdam, 1984) pp. 1–121.
- F. Omori, On the aftershocks of earthquakes, *J. Coll. Sci.* **7** (1895) 111.
- V. Pareto, *Cours d'Economie Politique*, (Rouge, Paris, 1897).
- K. Pearson, *The Grammar of Science*, 2nd ed. (Adam and Charles Black, London, 1900).

- T. C. Peterson and R. S. Vose, An overview of the global historical climatology network temperature database, *Bull. Am. Meteorol. Soc.* **78** (1997) 2837–2849.
- T. C. Peterson, R. Vose, R. Schmoyer and V. Razuvay, Global Historical Climatology Network (GHCN) quality control of monthly temperature data, *Int. J. Climatol.* **18** (1998a) 1169–1179.
- T. C. Peterson, T. R. Karl, P. F. Jamason, R. Knight, D. R. Easterling, The first difference method: maximizing station density for the calculation of long-term global temperature change, *Journal of Geophysical Research* **103** (1998b) 25967–25974.
- T. Phillips, The Mathematical Uncertainty Principle, *Monthly Essays on Mathematical Topics*, **November 2005**, American Mathematical Society. <http://www.ams.org/featurecolumn/archive/uncertainty.html>
- A. Rae, *Quantum-Physics: Illusion or Reality?* (Cambridge University Press, New York, 1988).
- L. F. Richardson, The problem of contiguity: An appendix to statistics of deadly quarrels, in *General Systems - Year Book of The Society for General Systems Research V*, eds. L. Von Bertalanffy and A. Rapoport, (Ann Arbor, MI, 1960), pp. 139-187.
- K. F. Riley, M. P. Hobson and S. J. Bence, *Mathematical Methods for Physics and Engineering*, 3rd ed. (Cambridge University Press, USA, 2006).
- T. Riste and D. Sherrington, eds., Spontaneous formation of space-time structures and criticality, Proc. NATO ASI, Geilo, Norway, (Kluwer, Dordrecht, 1991).
- C. Ruhla, *The Physics of Chance*, (Oxford University Press, 1992) p.217.
- M. Schroeder, *Fractals, Chaos and Power-laws*, (W. H. Freeman and Co., N. Y., 1990).
- A. M. Selvam, Deterministic chaos, fractals and quantumlike mechanics in atmospheric flows, *Can. J. Phys.* **68** (1990) 831-841. <http://xxx.lanl.gov/html/physics/0010046>
- A. M. Selvam, Universal quantification for deterministic chaos in dynamical systems, *Applied Math. Modelling* **17** (1993) 642-649. <http://xxx.lanl.gov/html/physics/0008010>
- A. M. Selvam and S. Fadnavis, Signatures of a universal spectrum for atmospheric inter-annual variability in some disparate climatic regimes, *Meteorology and Atmospheric Physics* **66** (1998) 87-112. <http://xxx.lanl.gov/abs/chao-dyn/9805028>
- A. M. Selvam and S. Fadnavis, Superstrings, cantorion-fractal spacetime and quantum-like chaos in atmospheric flows, *Chaos Solitons and Fractals* **10** (1999) 1321-1334. <http://xxx.lanl.gov/abs/chao-dyn/9806002>
- A. M. Selvam, D. Sen and S. M. S. Mody, Critical fluctuations in daily incidence of acute myocardial infarction, *Chaos, Solitons and Fractals* **11** (2000) 1175-1182. <http://xxx.lanl.gov/abs/chao-dyn/9810017>
- A. M. Selvam, Quantum-like chaos in prime number distribution and in turbulent fluid flows, *Apeiron* **8** (2001a) 29-64. <http://redshift.vif.com/JournalFiles/V08NO3PDF/V08N3SEL.PDF>
<http://xxx.lanl.gov/html/physics/0005067>
- A. M. Selvam, Signatures of quantum-like chaos in spacing intervals of non-trivial Riemann zeta zeros and in turbulent fluid flows, *Apeiron* **8** (2001b) 10-40. <http://redshift.vif.com/JournalFiles/V08NO4PDF/V08N4SEL.PDF> <http://xxx.lanl.gov/html/physics/0102028>
- A. M. Selvam, Cantorian fractal space-time fluctuations in turbulent fluid flows and the kinetic theory of gases, *Apeiron* **9** (2002a) 1-20. <http://redshift.vif.com/JournalFiles/V09NO2PDF/V09N2sel.PDF>
<http://xxx.lanl.gov/html/physics/9912035>
- A. M. Selvam, Quantumlike chaos in the frequency distributions of the bases A, C, G, T in Drosophila DNA, *Apeiron* **9** (2002b) 103-148. <http://redshift.vif.com/JournalFiles/V09NO4PDF/V09N4sel.pdf>
<http://arxiv.org/html/physics/0210068>
- A. M. Selvam, Quantumlike Chaos in the Frequency Distributions of the Bases A, C, G, T in human chromosome 1 DNA, *Apeiron* **11** (2004) 134-146. <http://redshift.vif.com/JournalFiles/V11NO3PDF/V11N3SEL.PDF>
<http://arxiv.org/html/physics/0211066>
- A. M. Selvam, A general systems theory for chaos, quantum mechanics and gravity for dynamical systems of all space-time scales, *ELECTROMAGNETIC PHENOMENA* **5**, No.2(15) (2005) 160-176. <http://arxiv.org/pdf/physics/0503028>
- A. M. Selvam, *Chaotic Climate Dynamics*, (Luniver Press, UK, 2007).
- D. Sornette, L. Knopoff, Y. Kagan and C. Vanneste, Rank-Ordering statistics of extreme events: application to the distribution of large earthquakes, arXiv:cond-mat/9510035v1, 6 Oct 1995.
- M. R. Spiegel, *Statistics*, Schaum's Outline Series in Mathematics, (McGraw-Hill, NY, 1961) pp.359.
- P. Steinhardt, Crazy crystals, *New Scientist* **25 Jan.** (1997) 32-35.
- A. A. Townsend, *The Structure of Turbulent Shear Flow*, 2nd ed. (Cambridge University Press, London, U.K., 1956) pp.115-130.
- G. K. Zipf, *Human Behavior and the Principle of Least Effort*, (Hafner, New York, 1949).